

HAMILTONIAN LINKING AND SYMPLECTIC PACKING

DYLAN CANT

ABSTRACT. In this short note, we make an observation relating symplectic packings of the standard symplectic ball by two sets and Hamiltonian linking.

1. Introduction

1.1. **Statement of results.** Denote by G be the group of compactly supported Hamiltonian diffeomorphisms of \mathbb{R}^{2n} , i.e., the group of time-1 maps of compactly supported Hamiltonian isotopies.

Let us say that two disjoint compact subsets $K_1, K_2 \subset \mathbb{R}^{2n}$ are *Hamiltonian unlinked* provided there is $\varphi \in G$ such that:

$$\varphi(K_1) \text{ and } \varphi(K_2) \text{ lie on opposite sides of a hyperplane.}$$

Otherwise we say they are *Hamiltonian linked*.

There is an obvious obstruction to being Hamiltonian unlinked, namely the two sets must be unlinkable via a compactly supported smooth isotopy.

Denote by $c(K_i)$ the *spectral capacity* of K_i , which is simply the Floer theory version¹ of Viterbo's generating function capacity introduced in [Vit92]; in this paper we follow the conventions of [AAC24, CZ24]. This is a normalized symplectic capacity. Then:

Theorem 1. *Suppose $K_1, K_2 \subset B(a)$ are compact disjoint subsets of the standard symplectic ball (of capacity $a > 0$). Either:*

- (1) K_1, K_2 are Hamiltonian linked,
- (2) $c(K_1) + c(K_2) \leq a$;

in other words, if two sets are Hamiltonian unlinked, then they obey a packing inequality; if two sets break the packing inequality, then they must be Hamiltonian linked.

Remark. The same result holds if $B(a)$ is replaced by a domain Ω satisfying $a = c(\Omega) = \gamma(\Omega)$, where $\gamma(\Omega)$ is the *spectral diameter* (i.e., the diameter of the space of isotopies supported in Ω with respect to Viterbo's spectral metric). In [AAC24] it is shown that $a = c(B(a)) = \gamma(B(a))$.

Corollary 2. *Let $K \subset \partial B(a)$ be a compact set with capacity $c(K) = b$ (there exist examples where K is a Lagrangian and $b = a/2$). Then the smaller ball $B(c) \subset B(a)$ is Hamiltonian linked with K if $c > a - b$. \square*

In dimension $n = 1$, this follows from smooth topology, since the only subsets $K \subset \partial B(a)$ with positive capacity are $K = \partial B(a)$. In higher dimensions $n > 1$,

Date: January 6, 2026.

¹Generating function capacities are known to equal Floer homology capacities; see [Vit96, Her04].

the sets K and $B(c)$ can be chosen to be smoothly unlinked, and so it produces a camel type theorem: *the ball $B(c)$ cannot be isotoped to $v + B(c)$ through balls of capacity c in the complement of K , if v is sufficiently large.*

Let us say that a compact domain $\Omega \subset \mathbb{C}^n$, with smooth and connected boundary $\partial\Omega$, is *boundary minimal* if:

$$c(\Omega \setminus U) < c(\Omega)$$

for all open sets U such that $U \cap \partial\Omega \neq \emptyset$. Then we recover a result of [Zil16] (in the case $\Omega = B(a)$); his argument used displacement energy while ours uses the packing versus linking theorem.

Corollary 3. *Any domain Ω satisfying $c(\Omega) = \gamma(\Omega)$ is boundary minimal.*

Proof of Corollary 3. If U is open and has non-empty intersection with $\partial\Omega$, then inside of $U \cap \Omega$ one can find a small ball which is unlinked with $\Omega \setminus U$. To find this ball, we use the assumption that $\partial\Omega$ is smooth and connected; it then follows each point $p \in \partial\Omega$ has a basis of neighborhoods of the form $(-\epsilon, \epsilon) \times D$ where:

- $\{0\} \times D$ is an open ball in $\partial\Omega$,
- $(-\epsilon, 0) \times D$ lies in the interior of Ω ,
- $(0, \epsilon) \times D$ lies in the non-compact connected component of $\mathbb{C}^n \setminus \Omega$.

Then, any point q in $(-\epsilon, 0) \times D$ can be moved by a Hamiltonian isotopy onto the opposite side of any hyperplane compared to Ω ; it follows that sufficiently small ball around q is unlinked with $\Omega \setminus U$, as desired. The result then follows from the packing inequality. \square

Remark. *It is interesting to wonder how the smoothness assumptions on $\partial\Omega$ can be weakened; see [BM25, §1.2] for related discussion.*

Boundary minimality is interesting as it implies the following closing-type lemma (see [Iri15, CDPT24, Xue22] for similar results):

Lemma 4. *Suppose that $\Omega \subset \mathbb{C}^n$ is a symplectically embedded Liouville domain which is boundary minimal. Let α be the intrinsic contact form on $\partial\Omega$ (using the Liouville domain structure of Ω). Then, for any non-empty open subset $V \subset \partial\Omega$ and any non-negative non-constant function f supported in V , there is a periodic Reeb orbit γ for the Reeb flow associated to $e^{-sf}\alpha$ which passes through V , for some $s \in (0, 1]$. Moreover:*

$$\text{period of } \gamma + \Gamma(\gamma) < c(\Omega)$$

where Γ is the cohomology class $[\lambda_{\mathbb{C}^n} - \lambda_\Omega] \in H_{\text{dR}}^1(\Omega)$. In particular, if the embedding is exact, one can bound the period of γ from above and conclude there is a periodic orbit of α with period $\leq c(\Omega)$ passing through any point of $\partial\Omega$.

This is proved in §2.5. As explained to the author by S. Matijević, the last sentence of the lemma, together with the results of [AK22, Iri22, AB23], implies the following characterization of Zoll domains amongst the strongly convex² domains.

Theorem 5. *A strongly convex domain Ω is Zoll if and only if it is boundary minimal.*

²Here we agree that *strongly convex* domains are those defined by equation $r = 1$ where r is a positive 1-homogeneous with everywhere positive definite Hessian (this is just a reformulation of any other reasonable definition of a “strongly” convex domain). This is an open subset of the space of convex domains, and this openness plays a role in the following proof.

Proof. If a convex domain with smooth boundary is boundary minimal, then Lemma 4, and the fact that $c(\Omega)$ is the shortest period of a periodic Reeb orbit (proved in [AK22, Iri22]) shows that Ω is Zoll. Here we do not need to assume that Ω is strongly convex.

Conversely, [AB23] proved that a uniformly convex Zoll domain is a local maximizer for the systolic ratio. Shrinking the domain slightly will decrease the volume, and so the systole must also decrease. Since small variations remain convex, it follows (again from [AK22, Iri22]) that $c(\Omega)$ decreases, as desired. \square

Our next result concerns the areas of J -holomorphic disks with boundary on Lagrangians. Here J is an ω -tame (and standard at infinity) almost complex structure (we call such a J *admissible*).

Corollary 6. *Let $K \subset \partial B(a)$ be a subset with capacity $b > 0$. Then each Lagrangian L in the interior of $B(a)$ either:*

- (1) *bounds a non-constant J -holomorphic disk with area at most $a - b$,*
- (2) *is linked with K ,*

where J is any admissible almost complex structure.

When $n = 1$ the result is trivial as every subset of $B(a)$ is smoothly linked with K , because the only subset with positive capacity is $K = \partial B(a)$. On the other hand, when $n > 1$, the result is symplectic in nature, as K and L are smoothly unlinked unless $K = \partial B(a)$. We prove Corollary 6 in §2.6.

Remark. Perhaps, when $n > 1$, every Lagrangian L in the interior of $B(a)$ is unlinked with some K with $c(K) \geq a/2$, in which case one proves the conjecture about the areas of disks being less than $a/2$; see §1.3.

Our next result discusses a case when sets are automatically unlinked.

Lemma 7. *If K_1 and K_2 are disjoint compact sets, each symplectomorphic to a starshaped set, then they are Hamiltonian unlinked.*

This is shown in §2.7. Theorem 1 then yields:

Corollary 8. *Suppose $a = c(\Omega) = \gamma(\Omega)$. Two disjoint sets $K_1, K_2 \subset \Omega$, each of which is symplectomorphic to a starshaped set, satisfy the packing inequality $c(K_1) + c(K_2) \leq a$. \square*

In fact, the proof of Theorem 1 shows that two disjoint sets K_1, K_2 which obey the maximum formula for spectral invariants also obey the packing inequality. It is known that the class of pairs of subsets of \mathbb{R}^{2n} which obey the maximum formula contains all pairs of the images of two embeddings of Liouville domains which are:

- (1) exact [HLRS16], or
- (2) π_1 -injective on the boundary [GT23],

We note that [HLRS16] also proves the the maximum formula for the generating function capacity for a pair of subsets of \mathbb{R}^{2n} which are separated by a hyperplane.

1.2. Discussion of proof. The key ingredients used to deduce Theorem 1 are:

- (1) The fact that the *spectral diameter of a ball equals its capacity*, i.e., $\gamma(B(a)) = c(B(a)) = a$, proved in [AAC24].
- (2) The maximum formula for spectral invariants, proved in [HLRS16, Tan22, GT23].

To deduce Corollary 6, one also appeals to the result in [Her04, CZ24].

1.3. Further questions.

1.3.1. *Areas of holomorphic disks.* Corollary 6 suggests the question:

Question 1. *Is every Lagrangian L in the interior of $B(a)$ Hamiltonian unlinked with some compact subset $K \subset \partial B(a)$ satisfying $c(K) \geq a/2$?*

As explained above, if the answer is “yes,” then it would solve the following conjecture (compare also with [CM18]):

Conjecture 2. *Every Lagrangian L in $B^{2n}(a)$ bounds non-constant J -holomorphic disks of area at most $a/2$, for all admissible J , if $n > 1$.*

1.3.2. *Minimal sets.* An interesting direction raised by Corollary 2 is that $\partial B(a)$ is a *minimal compact set* for the spectral capacity: any compact subset has a strictly smaller spectral capacity. Moreover, as explained above, the domain $B(a)$ is a boundary minimal Liouville domain, which implies some version of the closing lemma (see Lemma 4).

Question 3. *Which compact sets in \mathbb{R}^{2n} are minimal in the above sense? Which domains Ω are boundary minimal?*

It is not hard to see using [LS94] and [Her04, CZ24] that any compact connected Lagrangian submanifold is minimal, since removing a small ball will change its capacity from a positive number to zero. However, the boundary of a polydisk is never minimal.

Turning now to Zoll domains, Theorem 5 suggests the question:

Question 4. *Does every starshaped Zoll domain Ω satisfy $\gamma(\Omega) = c(\Omega)$? Is every starshaped domain satisfying $\gamma(\Omega) = c(\Omega)$ a Zoll domain?*

If the answer is “yes on both counts,” then $\gamma(\Omega) = c(\Omega)$ gives an alternative “spectral” characterization of when a starshaped domain is Zoll. Let us comment that Theorem 5 answers “yes” for the second part in the special case when Ω is known to be strictly convex.

It is known by the work of [GGM21, Mat24, Mat25, GRT25] that a starshaped domain is Zoll if and only if the first n Gutt-Hutchings capacities agree

$$c_1^{\text{GH}}(\Omega) = \dots = c_n^{\text{GH}}(\Omega)$$

(see [GH18a] for the definition). This raises a variant of Question 4:

Question 5. *Does every starshaped domain Ω with equal first and n th Gutt-Hutchings capacities satisfy the equality $\gamma(\Omega) = c(\Omega)$?*

Remark. *In the case $n = 2$, it is known from [GR25, GH18b] and [AAC24, §1.3] that $\gamma(\Omega) = c_2^{\text{GH}}(\Omega)$ holds whenever $\Omega \subset \mathbb{C}^2$ is a convex toric domain. Thus, if Ω is a Zoll convex toric domain, we conclude:*

$$\gamma(\Omega) = c_2^{\text{GH}}(\Omega) = c_1^{\text{GH}}(\Omega) = c(\Omega).$$

The middle inequality uses the characterization of Zoll convex domains, while the final inequality is well-known (at least for convex toric domains with smooth boundary, both $c_1^{\text{GH}}(\Omega)$ and $c(\Omega)$ equal the systole).

In fact more is true; as pointed out by the referee, the first part of Question 4 has a positive answer in the case $n = 2$ by appealing to [ABHS18, Proposition 4.3], which asserts that there is an ambient symplectomorphism of \mathbb{C}^2 taking any Zoll starshaped domain Ω to a rescaling of the standard closed ball.

In the case $n = 3$, one is tempted to answer “yes” to the first part of Question 4 by appealing to [CT25] which asserts that $\partial\Omega$ is strictly contactomorphic to $\partial B(a)$ for appropriate a ; however, whether this contactomorphism can be extended to a symplectomorphism of \mathbb{C}^3 taking Ω to $B(a)$ is not a priori clear — the argument in [ABHS18, Proposition 4.3] certainly depends on results specific to $n = 2$ (e.g., [ABHS18] use that $\text{Cont}(S^{2n-1})$ is path connected when $n = 2$ and such a result is unknown for higher values of n).

The notion of minimality also raises the question of how much the capacity drops when an open set is removed? On this topic, we prove the following:

Theorem 9. *If $0 < k < n$, and $0 \leq b_1 \leq b_2 \leq a$ then:*

$$K_{k,b_1,b_2} := \partial B(a) \cap \left\{ b_1 \leq \sum_{i=1}^k \pi |z_i|^2 \leq b_2 \right\}$$

satisfies $c(K_{k,b_1,b_2}) = \min\{b_2, a - b_1\}$.

In particular, even though the boundary of $B(a)$ is minimal, it contains proper compact subsets of any given smaller capacity. The proof is a simple application of Poisson commuting partitions of unity (i.e., using the toric structure) and embedding tricks, and is given in §2.8.

1.3.3. *Strong Arnol’d chord conjecture.* Boundary minimal sets are also related to the strong Arnol’d chord conjecture (see, e.g., [Zil16, Kan23, KZ24]). Indeed, adapting the argument of [Zil16], we can prove:

Theorem 10 (Ziltener). *Suppose that $\Omega \subset \mathbb{R}^{2n}$ is a boundary minimal exactly embedded Liouville domain, and $c(\Omega)$ is the minimal period of the Reeb flow on $\partial\Omega$ (e.g., if Ω is a uniformly convex Zoll domain). Let $\Lambda \subset \partial\Omega$ be a Legendrian. Then Λ has a Reeb chord of length at most $c(\Omega)/2$.*

Proof. Consider the map $\Lambda \times \mathbb{R}/c(\Omega)\mathbb{Z} \rightarrow \partial\Omega$ which evolves Λ by the Reeb flow. We have shown already in Lemma 4 that the Reeb flow on $\partial\Omega$ is $c(\Omega)$ -periodic, so the map is well-defined. If this map were injective, then it would produce an embedded Lagrangian $L \subset \partial\Omega$ with $\hbar(L) = c(\Omega)$, as in [Moh01, Zil16]. Then by [Her04, CZ24] this implies $c(L) = c(\Omega)$. However, $L \subset \partial\Omega$ is a proper compact subset of $\partial\Omega$, and so $c(L) < c(\Omega)$ by boundary minimality, thus the map could not have been an embedding. Thus there are two distinct points $x_1, x_2 \in \Lambda$ which lie on the same Reeb orbit, and this implies a Reeb chord of length $\leq c(\Omega)/2$. \square

Incidentally, using the projection $\pi : \partial B(1) \rightarrow \mathbb{C}P^{n-1}$, the fact that Lagrangians in $\partial B(1)$ (for $n > 1$) bound non-constant J -holomorphic disks of area strictly less than 1 implies any Lagrangian in $K \subset \mathbb{C}P^{n-1}$ bounds a disk of area strictly less than 1.³ Since K also bounds disks with symplectic area 1, it follows that K bounds a smooth disk with area less than 1/2. This disk can be lifted to $\pi^{-1}(K) \subset \partial B(1)$. Thus we conclude:

Corollary 11. *Every Lagrangian in $\partial B(1)$, with $n > 1$, bounds a smooth disk contained in $\partial B(1)$ with symplectic area at most 1/2.*

³Here we note that holomorphic disks of area less than 1 with boundary in $\partial B(1)$ miss the origin (by Gromov’s monotonicity theorem [Gro85]) and hence project to smooth disks in $\mathbb{C}P^{n-1}$ under the projection $\mathbb{C}^n \setminus \{0\} \rightarrow \mathbb{C}P^{n-1}$.

This result is certainly known to Ziltener (see [Zil16]), and presumably is known by other researchers as well. This is related to Conjecture 2, but does not imply it, as the disk is not a priori representable by a J -holomorphic disk for every admissible J ; moreover the argument applies only to Lagrangians contained in the boundary of the ball.

The *rationality constant* $\rho(L)$ of a Lagrangian L is infimal positive area of a smooth disk with boundary on L . Thus we have shown that $\rho(L) \leq a/2$ for all $L \subset \partial B(a)$ when $n > 1$. It is natural to compare with [DR15, Fai25] and ask the question:

Question 6. *Does every Lagrangian in $L \subset B(a)$ satisfy $\rho(L) \leq a/2$? If yes, are the Lagrangians L with $\rho(L) = a/2$ automatically contained in $\partial B(a)$?*

The question is only posed in \mathbb{R}^{2n} for $n > 1$, and the answer is known to be yes in both cases, in \mathbb{R}^4 , by the work of [CM18, DR15, Fai25].

Another consequence of the packing versus linking dichotomy is:

Theorem 12. *Suppose that $\Lambda_0, \Lambda_1 \subset \partial B(1)$ are disjoint and unlinked Legendrians, and let α denote the standard contact form on $\partial B(1)$. Then, for any $f \geq 0$, there exists a chord from $\Lambda_0 \cup \Lambda_1$ to $\Lambda_0 \cup \Lambda_1$ of length at most $1/2$ with respect to the contact form $e^{-f}\alpha$.*

Here two Legendrians in $\partial B(1)$ are unlinked provided there is a Legendrian isotopy of the link $\Lambda_0 \cup \Lambda_1$ sending the components to opposite sides of a hyperplane. We give the proof in §2.9. This result is interesting because it is stable with respect to a class of C^0 small perturbations of the ball (perturbations which push the ball inwards). It is known that the conclusion of the strong Arnol'd conjecture is not stable under C^0 small perturbations; see the discussion in [Kan23, KZ24].

1.3.4. Can zero capacity set link the ball?

Question 7. *If K links the ball $B(a)$, is it necessary that $c(K) > 0$?*

If the answer is positive, then one resolves the question of Ginzburg asking whether or not it holds that $c(N) > 0$ for all hypersurfaces N , see [Gin07, §3.3.4] and [CZ24, §1.4]. Unfortunately, one has:

Theorem 13. *For all sufficiently small $\epsilon > 0$, and $k < n$, the set:*

$$K = \partial B(a + \epsilon) \cap \left\{ \pi |z_i|^2 = 2\epsilon \text{ for all } i = 1, \dots, n-1 \right\}$$

links the ball and has capacity 2ϵ . Thus there are sets with arbitrarily small capacity which link the ball.

Proof. The claim about the capacity of K is well-known; indeed, the set is the elementary torus which has capacity 2ϵ provided ϵ is small enough compared to a . The linking theorem follows from the failure of the packing inequality between $B(a)$ and K in $B(a + \epsilon)$. \square

1.3.5. *Widths of domains.* Recently, [BT25] have established a general theorem which states that two domains V_1, V_2 in \mathbb{R}^{2n} satisfy a *max inequality* provided that there exist sufficiently large disjoint thickenings $V_i(r)$ defined for $r \in [0, 1]$. This means $V_i(r)$ is obtained by flowing V_i by a vector field which remains outwardly transverse to $\partial V_i(r)$ along the flow. Their theorem is expressed in terms the *width* of the thickening (see [BT25, Definition 1.2]).

Unfortunately, their theorem is stated only for non-negative Hamiltonians (our argument relies on both non-negative and non-positive Hamiltonians). More seriously, the max *inequality* is not sufficient for our argument to proceed (we require the max equality). This begs the question:

Question 8. *If V_1, V_2 are disjoint subsets of \mathbb{C}^n whose Buhovsky-Tanny widths are large enough, does it follow that $c(V_1) + c(V_2) \leq \gamma(V_1 \cup V_2)$?*

The thickenings considered in the above question should be disjoint.

A related concept to “width” is the maximal size of a disk cotangent bundle one can embed around a given Lagrangian L ; see [Vit90].

1.4. Acknowledgements. The author would like to acknowledge H. Alizadeh, M. S. Atallah, B. Bramham, O. Cornea, G. Dimitroglou-Rizell, Y. Ganor, V. Humilière, S. Matijević, L. Nakamura, S. Nemirovski, R. Leclercq, S. Seyfaddini, E. Shelukhin, C. Viterbo, and J. Zhang for useful discussions. The author also wishes to thank the anonymous referee for their insightful comments. The author is supported in his research by funding from the ANR project CoSy.

2. Proofs

We assume the reader is familiar with the set-up of spectral invariants and the spectral capacity, using the conventions of [AAC24, CZ24]. To each compactly supported Hamiltonian isotopy φ_t , which is generated by a compactly supported family of smooth functions H_t , one associates the spectral invariant $c(\varphi_t)$ of the unit element in Floer homology for H_t . For open sets U , one defines:

$$c(U) = \sup \{c(\varphi_t) : \varphi_t \text{ is supported in } U\},$$

and for compact sets K , one defines $c(K) = \inf \{c(U) : K \subset U \text{ with } U \text{ open}\}$.

2.1. Stability under Hamiltonian diffeomorphisms. One well-known fact that we will appeal to in our proof is:

Lemma 14. *Suppose that $g \in G$ and φ_t is a compactly supported isotopy. Then there is an equality of spectral invariants:*

$$c(g\varphi_tg^{-1}) = c(\varphi_t).$$

Proof. This can be proved using, e.g., a continuity argument and the fact that the action spectrum of $g_s\varphi_tg_s^{-1}$ is independent of s , where $g_s, s \in [0, 1]$, is a Hamiltonian isotopy generating g . See, e.g., [Vit92, Corollary 4.3] \square

2.2. Max formula. The version of the maximum formula we require is:

Lemma 15. *If $\varphi_{0,t}, \varphi_{1,t}$ are compactly supported Hamiltonian isotopies, and their supports lie on opposite sides of a hyperplane, then:*

$$c(\varphi_{0,t}\varphi_{1,t}) = \max \{c(\varphi_{0,t}), c(\varphi_{1,t})\}.$$

Proof. As explained in the introduction, this result is not new (and is proved in [HLRS16, GT23]), and is the *maximum formula for spectral invariants*. \square

2.3. The spectral diameter of the ball.

Lemma 16. *Suppose that φ_t is a Hamiltonian isotopy supported in $B(a)$. Then:*

$$\gamma(\varphi_t) = c(\varphi_t) + c(\varphi_t^{-1}) \leq a,$$

and the supremum of $\gamma(\varphi_t)$ over all such φ_t is a .

Proof. This is proved in [AAC24]. \square

2.4. Proof of Theorem 1. We can now prove the main theorem. Suppose that $K_1, K_2 \subset \Omega$ are disjoint compact sets, and $c(\Omega) = \gamma(\Omega) = a$. If they are Hamiltonian linked, then we are done. Therefore, assume they are Hamiltonian unlinked. Then there are disjoint open neighborhoods U_1, U_2 of K_1, K_2 which are also unlinked. Then:

$$c(K_1) + c(K_2) \leq c(U_1) + c(U_2) = \sup c(\varphi_{1,t}) + \sup c(\varphi_{2,t}),$$

where the supremums are over $\varphi_{i,t}$ supported in U_i . Now let $g \in G$ realize the unlinking of U_1, U_2 . Then §2.1 implies:

$$c(\varphi_{i,t}) = c(g\varphi_{i,t}g^{-1}) \text{ for } i = 1, 2.$$

On the other hand, since $g\varphi_{1,t}g^{-1}$ and $g\varphi_{2,t}g^{-1}$ are supported on opposite sides of a hyperplane, §2.1 and §2.2 imply:

$$c(\varphi_{1,t}\varphi_{2,t}^{-1}) = c(g\varphi_{1,t}\varphi_{2,t}^{-1}g^{-1}) = \max \{c(g\varphi_{1,t}g^{-1}), c(g\varphi_{2,t}^{-1}g^{-1})\}.$$

Similarly:

$$c(\varphi_{2,t}\varphi_{1,t}^{-1}) = c(g\varphi_{2,t}\varphi_{1,t}^{-1}g^{-1}) = \max \{c(g\varphi_{2,t}g^{-1}), c(g\varphi_{1,t}^{-1}g^{-1})\}.$$

Thus:

$$\begin{aligned} c(\varphi_{1,t}) + c(\varphi_{2,t}) &= c(g\varphi_{1,t}g^{-1}) + c(g\varphi_{2,t}g^{-1}) \\ &\leq c(\varphi_{1,t}\varphi_{2,t}^{-1}) + c(\varphi_{2,t}\varphi_{1,t}^{-1}) \leq a, \end{aligned}$$

where we have used $\gamma(\Omega) = a$ in the last line. Taking the supremum over $\varphi_{1,t}, \varphi_{2,t}$ completes the proof. \square

2.5. Proof of Lemma 4. The closing lemma we stated concerns symplectically embedding Liouville domains $\Omega \subset \mathbb{C}^n$ which are assumed to be boundary minimal (e.g., $\Omega = B(a)$).

Let $\Omega(sf) \subset \Omega$ be the subdomain obtained by flowing backwards by Ω 's intrinsic Liouville flow for time sf . Then $\lambda_\Omega|_{\partial\Omega(sf)} = e^{-sf}\alpha$.

It is well-known that the spectral capacity of $\Omega(sf) \subset \mathbb{C}^n$ is the period of some Reeb orbit γ_s for $e^{-sf}\alpha$, plus $\Gamma(\gamma_s)$ where $\Gamma = [\lambda_{\mathbb{C}^n} - \lambda_\Omega]$ is a cohomology class, i.e.,

$$\text{period of } \gamma_s + \Gamma(\gamma_s) = c(\Omega(sf)).$$

The set of periods of orbits which *do not* pass through V is independent of s , and the set of periods is closed and nowhere dense. In addition, the values of Γ are countable (when evaluated on elements of $H_1(\Omega, \mathbb{Z})$). Thus, if γ_s never passed through V , as s ranges over $(0, 1]$, then we would conclude that $c(\Omega(sf))$ was valued in a countable union of closed and nowhere dense sets. Since $s \mapsto c(\Omega(sf))$ is continuous (as is easy to prove by monotonicity and the definition of c via outer regularity) it follows that $c(\Omega(sf))$ is independent of s , contradicting boundary minimality of Ω . \square

2.6. Proof of Corollary 6. For this we only need to appeal to [Her04, Theorem 1.6] or [CZ24, Theorem 2], which relates the areas of holomorphic disks on Lagrangians to their spectral capacity. To show the statement is not vacuous, let us observe that one can take $K = \text{pr}^{-1}(\mathbb{R}\mathbb{P}^{n-1})$ where $\text{pr} : \partial B(a) \rightarrow \mathbb{C}\mathbb{P}^{n-1}$ is the Hopf fibration; see [Wei77, Aud88, Pol91] \square

2.7. Proof of Lemma 7. This is a straightforward isotopy extension type result (see, e.g., [MS12, §3]). Let us consider an embedding:

$$E = E_1 \sqcup E_2 : K_1 \sqcup K_2 \rightarrow \mathbb{R}^{2n}$$

where $K_i \subset \mathbb{R}^{2n}$ are compact starshaped sets (about the origin), and such that E_i extends to a symplectic map on a neighborhood of $K_i \subset \mathbb{R}^{2n}$. Define:

$$E_r = e^r E(e^{-r} z_1, e^{-r} z_2).$$

Then E_r is a symplectic isotopy, and $\omega(dE_r(-), \partial_r E_r)$ are closed one-forms on a neighborhood of K_i . Thus there is a smooth family of functions H_r such that:

$$E_r^* dH_r = \omega(dE_r(-), \partial_r E_r)$$

holds on a neighborhood of $K_1 \sqcup K_2$. It follows by the isotopy extension argument that the Hamiltonian flow by X_{H_r} on the interval $[0, s]$ takes E_0 to E_s .

The image of E_s is just a rescaled version of $E(e^{-s} K_1, e^{-s} K_2)$. The shrunken images $E_1(e^{-s} K_1)$ and $E_2(e^{-s} K_2)$ are eventually contained in disjoint balls around $E_1(0)$ and $E_2(0)$, and hence they can be separated by a hyperplane. Thus the rescaled images $e^s E_1(e^{-s} K_1)$ and $e^s E_2(e^{-s} K_2)$ are also separated by a hyperplane, as desired. \square

2.8. Proof of Theorem 9. Write $K = K_{k, b_1, b_2}$ and let K^c denote the complement of K inside of $\partial B(a)$. It suffices to prove the case $b_1 < b_2$.

Since $K \subset E(b_2, \dots, b_2, \infty, \dots)$ and $K \subset E(\infty, \dots, \infty, a - b_1, \dots)$, it follows that:

$$c(K) \leq \min \{a - b_1, b_2\}.$$

Now observe that K^c splits into two components:

- (1) $K_1^c = \left\{ \sum_{i=1}^k \pi |z_i|^2 \leq b_1 \right\} \cap \partial B(a)$,
- (2) $K_2^c = \left\{ \sum_{i=k+1}^n \pi |z_i|^2 \leq a - b_2 \right\} \cap \partial B(a)$.

We study the Hamiltonian orbit of K^c . For simplicity, suppose $b_1 \leq a - b_2$. First translate:

$$K_1^c \mapsto K_1^c + T e_n \text{ and } K_2^c \mapsto K_2^c,$$

for a large T . Then rotate by element of $g \in U(n)$ applied only to the translate of K_1^c , where g takes $T e_n$ to $T e_1$ and e_1 to $-e_n$; this rotation isotopy occurs in the complement of K_2^c provided T was large enough. Thus we see that the Hamiltonian orbit of K^c has a representative contained entirely in the region:

$$\left\{ \pi |z_n|^2 \leq a - b_2 \right\},$$

and hence $c(K^c) \leq a - b_2$. The other case $a - b_2 \leq b_1$ is similar, and we conclude $c(K^c) \leq \max \{a - b_2, b_1\}$.

Using the fact that K and K^c are Poisson commuting sets, we have:

$$a = c(K \cup K^c) \leq c(K) + c(K^c) \leq \min \{b_2, a - b_1\} + \max \{a - b_2, b_1\} = a,$$

and so all inequalities must be equalities. This proves:

$$c(K) = \min \{b_2, a - b_1\},$$

as desired. \square

2.9. Proof of Theorem 12. The construction of [Moh01] yields two disjoint Lagrangians inside $B(1)$ of capacities arbitrarily close to the minimal length of a Reeb chord of $\Lambda_0 \cup \Lambda_1$. These Lagrangians are unlinked because the Legendrians are unlinked. Thus the packing inequality holds, and we conclude the minimal length of a Reeb chord is at most $1/2$.

To see why the Lagrangians are unlinked, recall that they are defined as:

$$L_{i,R} := \Lambda_i \times \partial([0, R] \times [\delta, 1]) \rightarrow B(1),$$

where λ restricts to xdy where x, y are coordinates on the rectangle. The isotopy:

$$L_{i,s} := e^{s/2} L_{i,e^{-s}R}$$

is an exact isotopy — the area bounded by the loop remains constant. For s large enough, the projections of the Lagrangians to the contact ideal boundary enter arbitrarily small neighborhoods of the Legendrians. By assumption the Legendrians are unlinked, and so one can unlink the Lagrangians for s large enough via the symplectization lift of the isotopy unlinking Λ_i . \square

References

- [AAC24] H. Alizadeh, M. S. Atallah, and D. Cant, *The spectral diameter of a symplectic ellipsoid*, Accepted to J. Symplectic Geom., 2024.
- [AB23] A. Abbondandolo and G. Benedetti, *On the local systolic optimality of Zoll contact forms*, *Geom. Funct. Anal.* **33** (2023), no. 2, 299–363, doi:10.1007/s00039-023-00624-z. MR 4578460
- [ABHS18] A. Abbondandolo, B. Bramham, U. L. Hryniewicz, and P. A. S. Salomão, *Sharp systolic inequalities for Reeb flows on the three-sphere*, *Invent. Math.* **211** (2018), 687–778, doi:10.1007/s00222-017-0755-z.
- [AK22] A. Abbondandolo and J. Kang, *Symplectic homology of convex domains and Clarke’s duality*, *Duke Math. J.* **171** (2022), no. 3, 739–830, doi:10.1215/00127094-2021-0025. MR 4382979
- [Aud88] M. Audin, *Fibrés normaux d’immersions en dimension double, point doubles d’immersions lagrangiennes et plongements totalement réels*, *Comment. Math. Helv.* **63** (1988), 593–623.
- [BM25] Filip Bročić and Stefan Matijević, *A note on the wild symplectic ellipsoids*, arXiv:2512.21393, 2025.
- [BT25] L. Buhovsky and S. Tanny, *A local-to-global inequality for spectral invariants and an energy dichotomy for Floer trajectories*, *J. Fixed Point Theory Appl.* **27** (2025), no. 3, 1–37, doi:10.1007/s11784-024-01153-3. MR 4844096
- [CDPT24] J. Chaidez, I. Datta, R. Prasad, and S. Tanny, *Contact homology and higher dimensional closing lemmas*, *J. Mod. Dyn.* **20** (2024), 67–153, doi:10.3934/jmd.2024003.
- [CM18] K. Cieliebak and K. Mohnke, *Punctured holomorphic curves and Lagrangian embeddings*, *Invent. math.* **212** (2018), 213–295.
- [CT25] J. Chaidez and S. Tanny, *On zoll contact 5-spheres*, arXiv:2509.09639, 2025.
- [CZ24] D. Cant and J. Zhang, *On the spectral capacity of submanifolds*, arXiv:2409.14142, September 2024, Submitted.
- [DR15] G. Dimitroglou Rizell, *Uniqueness of extremal Lagrangian tori in the four-dimensional disc*, *Proceedings of the Gökova Geometry-Topology Conference* **22** (2015), 151–167.
- [Fai25] S. Faisal, *Extremal Lagrangian tori in toric domains*, arXiv:2504.13076, 04 2025, doi:10.48550/arXiv.2504.13076.
- [GGM21] V. L. Ginzburg, B. Z. Gürel, and M. Mazzucchelli, *On the spectral characterization of Besse and Zoll Reeb flows*, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **38** (2021), no. 3, 549–576, doi:10.1016/j.anihpc.2020.08.004. MR 4227045
- [GH18a] J. Gutt and M. Hutchings, *Symplectic capacities from positive S^1 -equivariant symplectic homology*, *Alg. Geom. Topol.* **18** (2018), 3537–3600.
- [GH18b] ———, *Symplectic capacities from positive S^1 -equivariant symplectic homology*, *Algebr. Geom. Topol.* **18** (2018), no. 6, 3537–3600, doi:10.2140/agt.2018.18.3537.
- [Gin07] V. L. Ginzburg, *Coisotropic intersections*, *Duke Math. J.* **140** (2007), no. 1, 111–163.

- [GR25] Jean Gutt and Vinicius G. B. Ramos, *Characterizing symplectic capacities on ellipsoids*, Rev. Mat. Iberoam. (2025), Published online first, doi:10.4171/RMI/1568.
- [Gro85] M. Gromov, *Pseudo holomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307–347.
- [GRT25] J. Gutt, V. G. B. Ramos, and S. Tanny, *Periodicity characterization by capacities for star-shaped domains*, arXiv:2511.18401, 2025.
- [GT23] Y. Ganor and S. Tanny, *Floer theory of disjointly supported Hamiltonians on symplectically aspherical manifolds*, Alg. Geom. Topol. **23** (2023), no. 2, 645–732.
- [Her04] D. Hermann, *Inner and outer Hamiltonian capacities*, Bull. Soc. math. France **132** (2004), no. 4, 509–541.
- [HLRS16] V. Humilière, F. Le Roux, and S. Seyfaddini, *Towards a dynamical interpretation of Hamiltonian spectral invariants on surfaces*, Geom. Topol. **20** (2016), no. 4, 2253–2334.
- [Iri15] K. Irie, *Dense existence of periodic Reeb orbits and ECH spectral invariants*, J. Mod. Dyn. **9** (2015), 357–363, doi:10.3934/jmd.2015.9.357.
- [Iri22] ———, *Symplectic homology of fiberwise convex sets and homology of loop spaces*, J. Symplectic Geom. **20** (2022), no. 2, 417–470, doi:10.4310/jsg.2022.v20.n2.a2.
- [Kan23] J. Kang, *On the strong Arnold chord conjecture for convex contact forms*, arXiv:2304.07016, 2023, doi:10.48550/arXiv.2304.07016.
- [KZ24] J. Kang and J. Zhang, *Strong Arnold chord conjecture via normalized capacities*, arXiv:2404.05150, 2024, doi:10.48550/arXiv.2404.05150.
- [LS94] F. Laudenbach and J.-C. Sikorav, *Hamiltonian disjunction and limits of Lagrangian submanifolds*, Int. Math. Res. Not. **1994** (1994), no. 4, 161–168.
- [Mat24] S. Matijević, *Positive (S^1 -equivariant) symplectic homology of convex domains, higher capacities, and Clarke’s duality*, arXiv:2410.13673v2, 10 2024, doi:10.48550/arXiv.2410.13673.
- [Mat25] ———, *Systolic S^1 -index and characterization of non-smooth Zoll convex bodies*, arXiv:2501.13856v2, 2 2025, doi:10.48550/arXiv.2501.13856.
- [Moh01] K. Mohnke, *Holomorphic disks and the chord conjecture*, Ann. Math. **154** (2001), 219–222, doi:10.2307/3062116.
- [MS12] D. McDuff and D. Salamon, *Introduction to symplectic topology*, 3rd ed., Oxford Graduate Texts in Mathematics, no. 27, Oxford University Press, 2012.
- [Pol91] L. Polterovich, *Monotone Lagrange submanifolds of linear spaces and the Maslov class in cotangent bundles*, Math. Z. **207** (1991), no. 2, 217–222.
- [Tan22] S. Tanny, *A max inequality for spectral invariants of disjointly supported Hamiltonians*, J. Symp. Geom. **20** (2022), no. 5, 1159–1213.
- [Vit90] C. Viterbo, *Plongements lagrangiens et capacités symplectiques de tores dans \mathbb{R}^{2n}* , C. R. Acad. Sci. Paris **311** (1990), 487–490.
- [Vit92] ———, *Symplectic topology as the geometry of generating functions*, Math. Ann. **292** (1992), no. 1, 685–710.
- [Vit96] ———, *Functors and computations in Floer homology with applications, part II*, arXiv:1805.01316v1, 1996.
- [Wei77] A. Weinstein, *Lectures on symplectic manifolds*, Regional Conference Series in Mathematics, vol. 29, American Mathematical Society, Providence, RI, 1977.
- [Xue22] J. Xue, *Closing lemma and KAM normal form*, arXiv:2207.06208, 2022, doi:10.48550/arXiv.2207.06208.
- [Zil16] F. Ziltener, *On the strict Arnold chord property and coisotropic submanifolds of complex projective space*, Int. Math. Res. Not. **2016** (2016), no. 3, 795–826, doi:10.1093/imrn/rnv153.

Email address: `dylan@dy lancant.ca`

DÉPARTAMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, PAVILLON ANDRÉ-AISENSTADT, 2920 CHEMIN DE LA TOUR, MONTRÉAL, QUÉBEC, H3T 1J4, CANADA